



# International Journal of Ayurveda Orientation

Volume 11, Issue 4, April 2026

---

## **FORMALIN EXPOSURE IN ANATOMY LABORATORIES: HEALTH RISKS AND INTEGRATED MODERN-AYURVEDIC PREVENTIVE STRATEGIES**

**Anop Singh , Dr. Gayatri Kalyanwat, Dr. Pankaj jain**

**Received:** April 14, 2026 | **Published:** April 20, 2026

---

Article ID: 150 | DOI: Pending  
[www.ijao.in](http://www.ijao.in)

# FORMALIN EXPOSURE IN ANATOMY LABORATORIES: HEALTH RISKS AND INTEGRATED MODERN-AYURVEDIC PREVENTIVE STRATEGIES

**Anop Singh , Dr. Gayatri Kalyanwat, Dr. Pankaj jain**

*Anop Singh (Siddheshwar Hanumanji Ayurveda College, Vaghada, Surendranagar, Gujarat.); Dr. Gayatri Kalyanwat (Govt. Ayurvedic Hospital, Ore, Sirohi, Rajasthan.); Dr. Pankaj jain (State Model Institute Of Ayurveda Science,Gandhinagar, Gujarat.)*

**Received:** Apr 14, 2026

**Published:** April 20, 2026

**ABSTRACT:** Formalin (aqueous formaldehyde solution) is ubiquitously employed in anatomy laboratories worldwide as a cadaveric preservative. Despite its practical utility, formalin poses substantial health hazards to medical students, faculty, and laboratory personnel through inhalation, dermal contact, and ocular exposure. Even at low concentrations of 0.1-2.7 ppm, formaldehyde causes significant mucous membrane irritation, with higher exposures associated with pulmonary edema, sensitization, and carcinogenesis [1,2,3]. The International Agency for Research on Cancer (IARC) and the U.S. National Institute for Occupational Safety and Health (NIOSH) classify formaldehyde as a probable human carcinogen, with no established safe threshold. This article reviews the physicochemical properties of formalin, documents observed health effects in anatomy dissection settings, outlines evidence-based modern preventive strategies including engineering controls, personal protective equipment (PPE), and administrative measures, and proposes an integrated approach incorporating validated Ayurvedic therapies — including Yashtimadhu (licorice), Anu Taila nasal oil, Nasyam therapy, and rasayana herbs — as adjunctive protective strategies [11,14,15,21]. The synthesis presented here aims to inform institutional policy and provide a holistic framework for minimizing occupational formalin-related morbidity in anatomy laboratory settings.

## INTRODUCTION

Gross anatomy dissection remains a cornerstone of medical education, providing students with



irreplaceable three-dimensional insight into human morphology. The preservation of cadavers for this purpose has historically relied on formalin — an aqueous solution of formaldehyde — owing to its potent antimicrobial and cross-linking properties. However, routine exposure to formalin in enclosed laboratory environments generates vapors that are now firmly established as hazardous to human health.

Studies investigating formaldehyde exposure in gross anatomy laboratories have documented vapour concentrations frequently exceeding recommended occupational exposure limits, with individual breathing-zone samples ranging from 0.43 to 2.01 ppm [1,2]. Students and staff in such environments report a spectrum of acute and chronic symptoms, and epidemiological evidence increasingly implicates long-term formaldehyde exposure in the development of nasopharyngeal cancer and leukaemia [3].

Despite the availability of modern engineering controls and PPE, compliance and infrastructure limitations — particularly in low- and middle-income countries — mean that many anatomy departments continue to expose trainees to unsafe formalin levels. There is thus a compelling need for cost-effective, complementary strategies. India's traditional system of medicine, Ayurveda, offers a rich pharmacopoeia of herbs and therapies documented for respiratory protection and immune modulation [11,17,19,20]. This review integrates modern occupational safety evidence with relevant Ayurvedic interventions to propose a dual-track preventive framework applicable to anatomy laboratories worldwide.

## Properties of Formalin

### Chemical and Physical Nature

Formalin is a 30–50% aqueous solution of formaldehyde (HCHO), stabilized with 10–15% methanol to prevent polymerization [9]. Formaldehyde itself is a colorless, flammable gas at room temperature with a characteristically pungent, suffocating odor detectable below 1 ppm. It has a molecular weight of 30.03 g/mol and a boiling point of  $-19^{\circ}\text{C}$ , meaning it exists as a vapor under standard laboratory conditions. Formaldehyde is heavier than air (vapor density 1.04 relative to air), causing vapors to accumulate at low-lying areas and floor levels in poorly ventilated rooms [2].

The compound is highly soluble in water and is readily absorbed by the respiratory mucosa, skin, and ocular surfaces. In anatomy laboratories, cadaveric embalming fluids typically contain formaldehyde at concentrations of approximately 3%, though commercial solutions as dilute as 0.625% have been documented in NIOSH health hazard evaluations [1]. Alternative preservation methods — including Thiel embalming and phenoxyethanol-based solutions — have been



proposed to reduce occupational formaldehyde exposure while preserving tissue quality adequate for dissection [9,10].

## Carcinogenic and Mutagenic Properties

Formaldehyde is classified as a Group 1 human carcinogen by the IARC and as a probable occupational carcinogen by NIOSH. Animal studies have demonstrated squamous cell carcinoma of the nasal passages in rats following inhalation exposure, and epidemiological investigations have linked chronic occupational exposure to elevated risks of nasopharyngeal cancer and myeloid leukaemia in humans [3]. Formaldehyde is genotoxic and mutagenic in multiple in vitro test systems, inducing DNA-protein cross-links, strand breaks, and chromosomal aberrations. Critically, no safe threshold of exposure for carcinogenic effects has been established, and NIOSH accordingly recommends reducing exposures to the lowest feasible level at all times.

## Sensitization Properties

Formalin is a potent chemical sensitizer. Repeated low-level exposure can induce IgE-mediated sensitization, following which even sub-threshold concentrations — well below 0.5 ppm — may trigger rhinitis, urticaria, asthma, and anaphylactoid reactions in sensitized individuals [3,6]. This sensitization phenotype is of particular concern in medical students, who face cumulative exposure across multiple dissection sessions over an academic year. Once sensitized, continued exposure perpetuates allergic disease and may necessitate removal from the laboratory environment.

## Observed Health Effects in Anatomy Laboratory Settings

### Respiratory and Mucous Membrane Effects

The respiratory tract is the primary target organ for formaldehyde toxicity. Inhalation of formalin vapors at concentrations as low as 0.1 ppm causes irritation of the nasal mucosa, pharynx, and larynx, manifesting as burning of the eyes, sneezing, nasal congestion, and coughing. At 0.3–2.7 ppm, sleep disturbance and worsening of pre-existing respiratory conditions have been documented [3]. Cross-sectional studies conducted among medical students attending gross anatomy dissection sessions have confirmed high prevalence rates of nasal discharge, throat irritation, watery eyes, and headache within the first weeks of cadaver exposure [4,5].



At higher concentrations of 10–20 ppm, more severe manifestations including chest tightness, palpitations, and bronchospasm are reported. Concentrations exceeding 50 ppm carry risk of pulmonary edema and chemical pneumonitis — life-threatening complications that necessitate emergency airway management. In susceptible individuals, particularly those with a history of atopy, even moderate vapor levels may precipitate acute asthma exacerbations [3,4].

## Dermatological and Ocular Effects

Skin contact with formalin solution causes both irritant contact dermatitis and, with repeated exposure, allergic contact dermatitis. The latter is immunologically mediated (Type IV hypersensitivity) and may persist or worsen despite reduced exposure intensity. Clinical features include erythema, pruritus, vesiculation, and fissuring of the hands and forearms — the most frequently affected sites in dissection laboratory workers [6,7]. Ocular exposure results in conjunctival irritation, excessive lacrimation, and — at high concentrations — corneal epithelial damage. Ayurvedic literature additionally documents the potential of topical formulations containing ingredients such as turmeric and neem for alleviating dermatitis associated with chemical sensitizers, though their specific efficacy against formalin-induced skin reactions requires formal clinical study [18].

## Allergic and Immunological Responses

A study assessing allergic responses among first-year medical students documented a significant incidence of formaldehyde-specific sensitization by the end of the first dissection semester, with skin prick testing and spirometry revealing both sensitization and subclinical bronchial hyperresponsiveness in a proportion of students [6]. These findings are concerning given that medical students represent a young, predominantly healthy cohort at the outset of a career involving potential lifelong formaldehyde exposure in clinical and research settings. Immunological studies further indicate that formaldehyde-hapten conjugates formed upon protein binding may serve as neoantigens driving chronic allergic disease.

## Systemic and Carcinogenic Risks

Systemic absorption of formaldehyde is generally limited by rapid metabolization via the formate pathway; however, inhalation of high concentrations causes systemic toxicity including metabolic acidosis and cardiovascular collapse. Ingestion — though unlikely in the laboratory context — is immediately life-threatening, with as little as 30 mL of a 37% solution reported as potentially fatal through corrosive gastrointestinal injury and multi-organ failure. Long-term low-level inhalation is epidemiologically associated with increased risk of leukaemia, nasal sinus cancer, and — in embalmers and anatomy professionals — elevated all-cause cancer mortality [3]. These



carcinogenic risks underscore the imperative for robust exposure control across all anatomy dissection environments.

## Modern Preventive Measures

The hierarchy of controls — elimination/substitution, engineering controls, administrative controls, and PPE — provides the evidence-based framework for managing occupational formaldehyde exposure in anatomy laboratories [7,8].

### Engineering Controls

Engineering controls represent the highest-order and most effective approach to formaldehyde hazard reduction. Key measures include:

- **General ventilation:** NIOSH recommends a minimum of 5-12 air changes per hour of fresh, unrecirculated air in gross anatomy laboratories, with some autopsy facility guidelines specifying up to 12 ACH [2,7].
- **Local exhaust ventilation (LEV):** Downdraft dissection tables with integrated exhaust have been demonstrated to significantly reduce cadaveric formaldehyde vapor levels at the breathing zone of students and demonstrators [2].
- **Substitution:** Use of alternative embalming fluids — including Thiel solution, phenoxyethanol-based preservatives, and glycerol-based formulations — can substantially reduce formaldehyde content while maintaining acceptable cadaveric quality for dissection. Reviews confirm the viability of these alternatives as lower-toxicity options [9,10].

### Administrative Controls

Administrative controls supplement engineering measures and include: formalized staff training programs on formalin hazards and safe handling procedures; limitation of individual dissection session duration and rotation of personnel in high-exposure tasks; pre-laboratory rinsing of cadavers with water to reduce surface formaldehyde concentration; scheduling of body cavity dissection (associated with higher vapor release) with adequate recovery intervals; implementation of a written Formaldehyde Exposure Control Plan as mandated by OSHA for workplaces with exposures at or above 0.1 ppm [7,8]; and establishment of biological and air monitoring programs to track exposure trends. Medical surveillance — including periodic respiratory function testing, skin examinations, and symptom questionnaires — should be offered to all regularly exposed personnel.



## Personal Protective Equipment (PPE)

When engineering and administrative controls cannot fully eliminate residual exposure risk, appropriate PPE is mandatory. OSHA and NIOSH specify the following minimum requirements for formalin handling in anatomy laboratories [7,8]:

- **Gloves:** Butyl rubber, neoprene, or nitrile gloves are recommended. Latex and polyvinyl chloride (PVC) gloves provide inadequate protection and must not be substituted.
- **Eye and face protection:** Chemical splash goggles or full face shields are required during direct handling of formalin solutions and during procedures likely to generate aerosols.
- **Respiratory protection:** Where vapour concentrations exceed the NIOSH REL of 0.016 ppm, half-face or full-face respirators with formaldehyde/organic vapor cartridges should be employed. At concentrations above 20 ppm (IDLH), supplied-air or self-contained breathing apparatus is required.

## Exposure Monitoring and Occupational Limits

Periodic personal breathing-zone air monitoring using validated methods (NIOSH Methods 3500/2016) is fundamental to ensuring that implemented controls achieve and maintain exposures below regulatory limits. OSHA's permissible exposure limit (PEL) is 0.75 ppm as an 8-hour TWA, with a 2 ppm short-term exposure limit (STEL). The NIOSH REL is far more stringent at 0.016 ppm (8-h TWA) reflecting carcinogenic risk, with a 15-minute ceiling of 0.1 ppm. The IDLH is 20 ppm [7,8]. Given that many anatomy laboratories in published studies exceed the NIOSH REL, institutions are strongly encouraged to target exposures as close to zero as technologically achievable.

## Ayurvedic Integration for Holistic Protection

While modern engineering and PPE-based controls address the primary occupational hazard, Ayurveda — the ancient Indian system of medicine — offers a complementary pharmacological and therapeutic framework targeting respiratory resilience, mucosal integrity, immune modulation, and detoxification. The following Ayurvedic interventions are supported by traditional textual references and, where available, emerging pharmacological evidence. It must be emphasized that these measures **supplement** — and must never replace — CDC/NIOSH-recommended primary controls.



## Yashtimadhu (*Glycyrrhiza glabra* – Licorice Root)

Yashtimadhu (licorice root) is one of the most extensively documented herbs in the Ayurvedic pharmacopoeia for upper respiratory protection. Its active constituents — glycyrrhizin, liquiritin, and isoliquiritin — have demonstrated anti-inflammatory, antioxidant, antiviral, and expectorant properties in experimental models [21]. Classical Ayurvedic texts describe Yashtimadhu as a *kanthashodhana* (throat-cleansing) and *kasahara* (antitussive) drug, with particular utility in pharyngitis, tonsillitis, and allergic cough — symptoms consistent with formalin-induced mucous membrane irritation [11,12,13]. In the context of anatomy laboratory exposure, *Yashtimadhu Ghanvati* (tablet formulation) or decoction taken before and after dissection sessions may offer mucoprotective and anti-inflammatory benefit to the upper airway. Its immunomodulatory properties may additionally attenuate the sensitization process triggered by repeated formaldehyde exposure.

## Anu Taila and Nasya (Nasal Oil Therapy)

Anu Taila is a classical Ayurvedic polyherbal nasal oil formulation documented for the management of sinusitis, nasal allergy, and upper respiratory inflammation [14]. Administered via *Nasyam* — the Ayurvedic para-nasal instillation of medicated oils — it is believed to form a protective lipid coating on the nasal mucosa, potentially reducing direct formaldehyde-mucosal contact and attenuating local inflammatory responses [15]. *Nasyam* is classified in Ayurveda as one of the Panchakarma (five detoxification procedures) and is specifically indicated for diseases of the head, neck, and upper respiratory tract. Pre-dissection nasal oil application could theoretically serve as a low-cost mucosal barrier measure, though rigorous clinical trials in the context of chemical exposure are yet to be conducted.

## Rasayana Herbs for Immune Modulation

Ayurvedic rasayana (rejuvenative) therapy encompasses a class of herbs and formulations designed to enhance *ojas* (vitality and immune resilience). Several such herbs are particularly relevant to formalin-exposed individuals:

- **Turmeric (*Curcuma longa*):** Curcumin, the principal active constituent, exhibits potent anti-inflammatory and antioxidant properties and has been shown to modulate NF-κB-mediated inflammatory pathways implicated in chemical-induced airway injury. Its incorporation into daily diet or as a supplement may reduce systemic inflammatory burden in exposed individuals [17,19].
- **Tulsi (*Ocimum sanctum*):** Sacred basil is documented in Ayurveda as a *shwasahara* (anti-



dyspnoeic) and immunoadaptogenic herb. Clinical and experimental studies support its bronchodilatory and mast cell stabilizing properties, which may be beneficial in mitigating formaldehyde-induced bronchospasm and airway hypersensitivity [19,20].

- **Ginger (*Zingiber officinale*):** Ginger's gingerols and shogaols possess anti-inflammatory and antihistaminic properties. In Ayurvedic practice, ginger-based decoctions are routinely prescribed for respiratory allergies associated with pollution or chemical irritants — a context analogous to formalin exposure [17].

## Snehana (Oleation Therapy) and General Detoxification

Snehana, or Ayurvedic oleation therapy, involves the systemic or topical application of medicated oils or ghee to nourish body tissues, facilitate the mobilization of lipid-soluble toxins, and prepare the body for deeper detoxification procedures (shodhana) [16]. Given that formaldehyde and its metabolites (particularly formate) are water-soluble and rapidly excreted, classical Ayurvedic detoxification using specific herbal decoctions and dietary modifications may support hepatic metabolic clearance pathways. Regular Abhyanga (full-body oil massage) using sesame or medicated oils is also advocated in Ayurveda for stress reduction, skin barrier fortification, and general immune enhancement — all potentially beneficial for laboratory-exposed students.

## Steam Inhalation and Herbal Fumigation

Herbal steam inhalation with formulations containing eucalyptus, camphor, or tulsi is a traditional Ayurvedic recommendation for clearing upper respiratory congestion and mucosal irritation caused by air pollutants and chemical vapors [17]. Employed post-exposure as a decontamination ritual, steam inhalation may help clear residual formaldehyde adsorbed to nasal and pharyngeal mucosae, while the anti-inflammatory and antimicrobial properties of dissolved herbal constituents may reduce secondary mucosal inflammation. Similarly, Ayurvedic approaches to strengthening mucosal immunity — including probiotics and fermented herbal preparations — have been described as adjuncts in allergy management [20].

## Proposed Integrated Preventive Framework

An integrated framework for formalin exposure management in anatomy laboratories should operate across three tiers: (1) primary institutional controls following the CDC/NIOSH/OSHA hierarchy of controls; (2) individual-level PPE and safe work practices; and (3) adjunctive Ayurvedic wellness measures supporting mucosal integrity, immune function, and respiratory resilience. This tiered approach recognizes that no single intervention is sufficient and that comprehensive protection requires simultaneous action at organizational and individual levels.



Institutions in South Asian contexts — where both Ayurvedic expertise and infrastructure limitations frequently coexist — are particularly well-positioned to adopt this integrated approach. Implementation of downdraft tables and improved ventilation should remain the priority investment; Ayurvedic adjuncts such as *Yashtimadhu Ghanvati*, Anu Taila nasal application, and rasayana herbal supplementation represent low-cost, culturally acceptable, and pharmacologically plausible additions to the standard protective protocol [11,14,21].

## CONCLUSION

Formalin exposure in anatomy laboratories constitutes a significant and preventable occupational health hazard, with well-documented acute irritative, sensitizing, and long-term carcinogenic consequences. The evidence base supporting modern engineering controls — particularly adequate ventilation, local exhaust at the dissection site, and substitution with lower-formaldehyde preservatives — is robust and should inform institutional policy universally. PPE and structured administrative controls provide critical additional protection layers.

Ayurvedic integrative strategies — including *Yashtimadhu* for mucosal protection, Nasyam nasal therapy for upper airway defense, and rasayana herbs for immune modulation — offer a culturally rooted, pharmacologically plausible complement to conventional occupational safety measures. Future research should prioritize controlled clinical trials evaluating the efficacy of these Ayurvedic interventions in documented formaldehyde-exposed populations, with standardized outcome measures for respiratory sensitization, inflammatory biomarkers, and quality of life. Until such evidence accumulates, Ayurvedic measures should be transparently positioned as adjuncts — never substitutes — to primary engineering and PPE controls, administered under qualified medical supervision.

## REFERENCES

- [1] Estimation of occupational formaldehyde exposure in cadaver dissection laboratories. PMC6952696. <https://pmc.ncbi.nlm.nih.gov/articles/PMC6952696/>
- [2] Formaldehyde Exposures in a Gross Anatomy Laboratory. CDC/MMWR. <https://www.cdc.gov/mmWR/preview/mmwrhtml/00001222.htm>
- [3] Health Effects of Low-Dose Formaldehyde. PMC12273900.



<https://pmc.ncbi.nlm.nih.gov/articles/PMC12273900/>

[4] Effect of Formalin Exposure on Medical Students. Journal of Pharmacy and Therapeutics. <https://jptcp.com/index.php/jptcp/article/download/11133/10473/22428>

[5] Effects of Formalin Exposure Among Students Attending Anatomy Dissection in Medical College: A Cross-Sectional Study. CPD Clinical Biochemistry. <https://cpdclinicalbiochem.co.uk/article/effects-of-formalin-exposure-among-students-attending-anatomy-dissection-in-medical-college-a-cross-sectional-study-592/>

[6] A Study of Allergic Responses of Formaldehyde in First Year Medical Students. International Journal of Research in Medical Sciences. <https://www.msjonline.org/index.php/ijrms/article/download/8569/5946>

[7] Formaldehyde Exposure Control Plan. University of Wisconsin-Madison Environmental Health & Safety. <https://ehs.wisc.edu/wp-content/uploads/sites/1408/2020/09/EHS-EOH-GUI-006-V01.pdf>

[8] OSHA Fact Sheet: Formaldehyde. <https://www.odu.edu/sites/default/files/documents/formaldehyde.pdf>

[9] Alternative Natural and Chemical Substances to Traditional Formalin-Based Embalming Fluid for Cadaveric Dissection: A Review. Indian Journal of Clinical Anatomy and Physiology. <https://www.ijcap.org/article-details/19220>

[10] Thiel Embalming Method Used for Anatomy Dissection as an Alternative to Standard Formalin Embalming. Folia Morphologica. [https://journals.viamedica.pl/fovia\\_morphologica/article/view/89320](https://journals.viamedica.pl/fovia_morphologica/article/view/89320)

[11] Yashtimadhu Churna: Health Benefits, Medicinal Uses, Dosage, Side Effects and Precautions. Netmeds Health Library. <https://www.netmeds.com/c/health-library/post/yashtimadhu-churna-health-benefits-medicinal-uses-method-dosage-side-effects-and-precautions>

[12] Nagarjun Yashtimadhu Ghanvati: Uses in Dry, Allergic and Wet Cough, Pharyngitis, Tonsillitis. ShopHealthy. <https://www.shophealthy.in/nagarjun-yashtimadhu-ghanvati-1200tab>

[13] Yashtimadhu Ghanvati. Desi Dawaiya. <http://desidawaiya.com/shop/vati-gutika/yashtimadhu-ghanvati/>

[14] Anu Taila: Ayurvedic Oil for Sinusitis and Nasal Allergy Relief. AyurvedaBay. <https://www.ayurvedabay.com/anu-taila.html>



- [15] Nasyam Ayurvedic Treatment. Sreedhareeyam Ayurvedic Medical College and Hospital. <https://www.sreedhareeyam.com/hospital/blog/nasyam-ayurvedic-treatment-132>
- [16] Ayurvedic Oleation and Snehana for Body Tissue Nourishment and Detox. Fazlani Nature's Nest. <https://fazlaninaturesnest.com/snehana-oleation-ayurveda-therapy-what-is-it-types-and-benefits-it-offers-to-body-tissues/>
- [17] Pollution Causing Cough and Cold? Try Ayurvedic Home Remedies. Maharishi Ayurveda India. <https://maharishiayurvedaindia.com/blogs/wellness-need/ayurvedic-remedies-for-pollution-related-cough-and-cold>
- [18] Ayurvedic Ingredients for Sensitive Skin. VedicDerm. <https://vedicderm.com/hi/blogs/news/ayurvedic-ingredients-for-sensitive-skin>
- [19] Top 15 Ayurvedic Remedies for Air Pollution Issues. Tigris Valley. <https://tigrisvalley.com/ayurvedic-remedies-for-air-pollution-issues/>
- [20] Ayurvedic Approach to Strengthening Immunity Against Allergies. Ishani Ayurveda. <https://www.ishaniayurveda.com/blog/ayurvedic-approach-to-strengthening-immunity-against-allergies/>
- [21] Pharmacological Profile of Yashtimadhu in Urdhwajatru Vikaras. Journal of Ayurveda and Integrative Medicine Sciences. <https://jaims.in/jaims/article/download/4513/7755?inline=1>

#### HOW TO CITE THIS ARTICLE:

**APA:** Anop Singh , Dr. Gayatri Kalyanwat, Dr. Pankaj jain (2026). Formalin Exposure in Anatomy Laboratories: Health Risks and Integrated Modern-Ayurvedic Preventive Strategies. *International Journal of Ayurveda Orientation*, 11(4), 21-31.

**MLA:** Anop Singh , et al. "Formalin Exposure in Anatomy Laboratories: Health Risks and Integrated Modern-Ayurvedic Preventive Strategies." *International Journal of Ayurveda Orientation* 11.4 (2026): 21-31.

**Vancouver:** Anop et al. Formalin Exposure in Anatomy Laboratories: Health Risks and Integrated Modern-Ayurvedic Preventive Strategies. *International Journal of Ayurveda Orientation* 2026;11(4):21-31.